

Latest Heat Transfer

1. Unit of thermal conductivity in M.K.S. units is

- (a) kcal/kg m² °C
- (b) kcal-m/hr m² °C
- (c) kcal/hr m² °C
- (d) kcal-m/hr °C
- (e) kcal-m/m² °C.

Ans: b

2. Unit of thermal conductivity in S.I. units is

- (a) J/m² sec
- (b) J/m °K sec
- (c) W/m °K
- (d) (a) and (c) above
- (e) (b) and (c) above.

Ans: e

3. Thermal conductivity of solid metals with rise in temperature normally

- (a) increases
- (b) decreases
- (c) remains constant
- (d) may increase or decrease depending on temperature
- (e) unpredictable.

Ans: b

4. Thermal conductivity of non-metallic amorphous solids with decrease in temperature

- (a) increases
- (b) decreases
- (c) remains constant
- (d) may increase or decrease depending on temperature
- (e) unpredictable.

Ans: b

5. Heat transfer takes place as per -

- (a) zeroth law of thermodynamics
- (b) first law of thermodynamic
- (c) second law of the thermodynamics
- (d) Kirchoff's law (e) Stefan's law.

Ans: c

6. When heat is transferred from one particle of hot body to another by actual motion of the heated particles, it is referred to as heat transfer by

- (a) conduction
- (b) convection
- (c) radiation
- (d) conduction and convection
- (e) convection and radiation.

Ans: a

7. When heat is transferred from hot body to cold body, in a straight line, without affecting the intervening medium, it is referred as heat transfer by

- (a) conduction
- (b) convection
- (c) radiation
- (d) conduction and convection
- (e) convection and radiation.

Ans: c

8. Sensible heat is the heat required to

- (a) change vapour into liquid
- (b) change liquid into vapour
- (c) increase the temperature of a liquid or vapour
- (d) convert water into steam and superheat it
- (e) convert saturated steam into dry steam.

Ans: c

9. The insulation ability of an insulator with the presence of moisture would

- (a) increase
- (b) decrease
- (c) remain unaffected
- (d) may increase/decrease depending on temperature and thickness of insulation
- (e) none of the above.

Ans: b

10. When heat is transferred by molecular collision, it is referred to as heat transfer by

- (a) conduction
- (b) convection
- (c) radiation
- (d) scattering
- (e) convection and radiation.

Ans: b

11. Heat transfer in liquid and gases takes place by

- (a) conduction
- (b) convection
- (c) radiation
- (d) conduction and convection
- (e) convection and radiation.

Ans: b

12. Which of the following is the case of heat transfer by radiation

- (a) blast furnace
- (b) heating of building
- (c) cooling of parts in furnace
- (d) heat received by a person from fireplace
- (e) all of the above.

Ans: d

13. Heat is closely related with

- (a) liquids
- (b) energy
- (c) temperature
- (d) entropy
- (e) enthalpy.

Ans: c

14. Pick up the wrong case. Heat flowing from one side to other depends directly on

- (a) face area
- (b) time
- (c) thickness
- (d) temperature difference
- (e) thermal conductivity.

Ans: c

15. Metals are good conductors of heat because

- (a) their atoms collide frequently
- (b) their atoms-are relatively far apart
- (c) they contain free electrons
- (d) they have high density
- (e) all of the above.

Ans: a

16. Which of the following is a case of steady state heat transfer

- (a) I.C. engine
- (b) air preheaters
- (c) heating of building in winter
- (d) all of the above
- (e) none of the above.

Ans: e

17. Total heat is the heat required to

- (a) change vapour into liquid
- (b) change liquid into vapour
- (c) increase the temperature of a liquid or vapour
- (d) convert water into steam and superheat it
- (e) convert saturated steam into dry steam.

Ans: d

18. Cork is a good insulator because it has

- (a) free electrons
- (b) atoms colliding frequency
- (c) low density
- (d) porous body
- (e) all of the above.

Ans: d

19. Thermal conductivity of water in general with rise in temperature

- (a) increases
- (b) decreases
- (c) remains constant
- (d) may increase or decrease depending on temperature
- (e) none of the above.

Ans: d

20. Thermal conductivity of water at 20°C is of the order of

- (a) 0.1
- (b) 0.23
- (c) 0.42
- (d) 0.51
- (e) 0.64.

Ans: d

21. Temperature of steam at around 540°C can be measured by

- (a) thermometer
- (b) radiatiouv pyrometer
- (c) thermistor
- (d) thermocouple
- (e) thermopile.

Ans: d

22. Thermal conductivity of air at room temperature in kcal/m hr °C is of the order of

- (a) 0.002
- (b) 0.02
- (c) 0.01
- (d) 0.1
- (e) 0.5.

Ans: b

23. The time constant of a thermocouple is

- (a) the time taken to attain the final temperature to be measured
- (b) the time taken to attain 50% of the value of initial temperature difference
- (c) the time taken to attain 63.2% of the value of initial temperature difference
- (d) determined by the time taken to reach 100°C from 0°C
- (e) none of the above.

Ans: c

24. Thermal conductivity of air with rise in temperature

- (a) increases
- (b) decreases
- (c) remains constant
- (d) may increase or decrease depending on temperature
- (e) none of the above.

Ans: a

25. Heat flows from one body to other when they have

- (a) different heat contents
- (b) different specific heat

- (c) different atomic structure
- (d) different temperatures
- (e) none of the above.

Ans: d

26. The concept of overall coefficient of heat transfer is used in heat transfer problems of

- (a) conduction
- (b) convection
- (c) radiation
- (d) all the three combined
- (e) conduction and convection.

Ans: e

27. In heat transfer, conductance equals conductivity (kcal/hr/sqm/°C/cm) divided by

- (a) hr (time)
- (b) sqm (area)
- (c) °C (temperature)
- (d) cm (thickness)
- (e) kcal (heat).

Ans: d

28. The amount of heat flow through a body by conduction is

- (a) directly proportional to the surface area of the body
- (b) directly proportional to the temperature difference on the two faces of the body
- (c) dependent upon the material of the body
- (d) inversely proportional to the thickness of the body
- (e) all of the above.

Ans: e

29. Which of the following has least value of conductivity

- (a) glass
- (b) water
- (c) plastic
- (d) rubber
- (e) air.

Ans: e

30. Which of the following is expected to have highest thermal conductivity

- (a) steam
- (b) solid ice
- (c) melting ice
- (d) water
- (e) boiling water.

Ans: b

6-31. Thermal conductivity of glass-wool varies from sample to sample because of variation in

- (a) composition
- (b) density
- (c) porosity

- (d) structure
- (e) all of the above.

Ans: e

32. Thermal conductivity of a material may be defined as the

- (a) quantity of heat flowing in one second through one cm cube of material when opposite faces are maintained at a temperature difference of 1°C
- (b) quantity of heat flowing in one second through a slab of the material of area one cm square, thickness 1 cm when its faces differ in temperature by 1°C
- (c) heat conducted in unit time across unit area through unit thickness when a temperature difference of unity is maintained between opposite faces
- (d) all of the above
- (e) none of the above.

Ans: d

33. Which of the following has maximum value of thermal conductivity

- (a) aluminium
- (b) steel
- (c) brass
- (d) copper
- (e) lead.

Ans: a

34. Moisture would find its way into insulation by vapour pressure unless it is prevented by

- (a) high thickness of insulation
- (b) high vapour pressure
- (c) less thermal conductivity insulator
- (d) a vapour seal
- (e) all of the above.

Ans: d

35. Heat is transferred by all three modes of transfer, viz, conduction, convection and radiation in

- (a) electric heater
- (b) steam condenser
- (c) melting of ice
- (d) refrigerator condenser coils
- (e) boiler.

Ans: e

36. According to Prevost theory of heat exchange

- (a) it is impossible to transfer heat from low temperature source to a high temperature source
- (b) heat transfer by radiation requires no medium
- (c) all bodies above absolute zero emit radiation
- (d) heat transfer in most of the cases takes place by combination of conduction, convection and radiation
- (e) rate of heat transfer depends on thermal conductivity and temperature difference.

Ans: c

37. The ratio of heat flow Q_1/Q_2 from two walls of same thickness having their thermal conductivities as $AT_j - 2K_2$ will be

- (a) 1
- (b) 0.5
- (c) 2
- (d) 0.25
- (e) 4.0

Ans: c

38. Heat transfer by radiation mainly depends upon

- (a) its temperature
- (b) nature of the body
- (c) kind and extent of its surface
- (d) all of the above
- (e) none of the above.

Ans: d

39. Thermal diffusivity is

- (a) a dimensionless parameter
- (b) function of temperature
- (c) used as mathematical model
- (d) a physical property of the material
- (e) useful in case of heat transfer by radiation.

Ans: d

40. Thermal diffusivity of a substance is .

- (a) proportional of thermal conductivity
- (b) inversely proportional to k
- (c) proportional to (k)
- (d) inversely proportional to k^2
- (e) none of the above.

Ans: a

41. Unit of thermal diffusivity is

- (a) m^2/hr
- (b) $m^2/hr^\circ C$
- (c) $kcal/m^2 hr$
- (d) $kcal/m.hr^\circ C$
- (e) $kcal/m^2 hr^\circ C$.

Ans: a

43. Thermal conductivity of wood depends on

- (a) moisture
- (b) density
- (c) temperature
- (d) all of the above
- (e) none of the above.

Ans: d

44. In convection heat transfer from hot flue gases to water tube, even though flow may be turbulent, a laminar flow region (boundary layer of film) exists close to the tube. The heat transfer through this film takes place by

- (a) convection
- (b) radiation
- (c) conduction
- (d) both convection and conduction
- (e) none of the above.

Ans: c

45. Film coefficient is defined as

- (a) Inside diameter of tube
- (b) Equivalent thickness of film
- (c) Thermal conductivity / Equivalent thickness of film
- (d) Thermal conductivity / Molecular diffusivity of momentum
- (e) Thermal diffusivity
- (f) Film coefficient x Inside diameter
- (g) Thermal conductivity
- (h) none of the above.

Ans: b

46. Heat conducted through unit area and unit thick face per unit time when temperature difference between opposite faces is unity, is called

- (a) thermal resistance
- (b) thermal coefficient
- (c) temperature gradient
- (d) thermal conductivity
- (e) heat-transfer.

Ans: d

49. The rate of energy emission from unit surface area through unit solid angle, along a normal to the surface, is known as

- (a) emissivity
- (b) transmissivity
- (c) reflectivity
- (d) intensity of radiation
- (e) absorptivity.

Ans: d

50. Emissivity of a white polished body in comparison to a black body is

- (a) higher
- (b) lower
- (c) same
- (d) depends upon the shape of body
- (e) none of the above.

Ans: b

51. A grey body is one whose absorptivity

- (a) varies with temperature
- (b) varies with wavelength of the incident ray
- (c) is equal to its emissivity
- (d) does not vary with temperature and wavelength of the incident ray
- (e) none of the above.

Ans: c

53. Two balls of same material and finish have their diameters in the ratio of 2 : 1 and both are heated to same temperature and allowed to cool by radiation. Rate of cooling by big ball as compared to smaller one will be in the ratio of

- (a) 1 : 1
- (b) 2 : 1
- (c) 1 : 2
- (d) 4 : 1
- (e) 1 : 4.

Ans: c

55. A non-dimensional number generally associated with natural convection heat transfer is

- (a) Grashoff number
- (b) Nusselt number
- (c) Weber number
- (d) Prandtl number
- (e) Reynold number.

Ans: a

56. LMTD in case of counter flow heat exchanger as compared-to parallel flow heat exchanger is

- (a) higher
- (b) lower
- (c) same
- (d) depends on the area of heat exchanger
- (e) depends on temperature conditions.

Ans: a

57. In heat exchangers, degree of approach is defined as the difference between temperatures of

- (a) cold water inlet and outlet
- (b) hot medium inlet and outlet
- (c) hot medium outlet and cold water inlet
- (d) hot medium outlet and cold water outlet
- (e) none of the above.

Ans: d

58. In counter flow heat exchangers

- (a) both the fluids at inlet (of heat ex-changer where hot fluid enters) are in their coldest state
- (b) both the fluids at inlet are in their hot-test state
- (c) both the fluids .at exit are in their hottest state
- (d) one fluid is in hottest state and other in coldest state at inlet
- (e) any combination is possible depending on design of heat exchanger.

Ans: b

59. A steam pipe is to be insulated by two insulating materials put over each other. For best results

- (a) better insulation should be put over pipe and better one over it
- (b) inferior insulation should be put over pipe and better one over it
- (c) both may be put in any order
- (d) whether to put inferior OIL over pipe or the better one would depend on steam temperature
- (e) unpredictable.

Ans: a

61. Fourier's law of heat conduction is valid for

- (a) one dimensional cases only
- (b) two dimensional cases only
- (c) three dimensional cases only
- (d) regular surfaces having non-uniform temperature gradients
- (e) irregular surfaces.

Ans: a

62. According of Kirchhoff's law,

- (a) radiant heat is proportional to fourth power of absolute temperature
- (b) emissive power depends on temperature
- (c) emissive power and absorptivity are constant for all bodies
- (d) ratio of emissive power to absorptive power is maximum for perfectly black body
- (e) ratio of emissive power to absorptive power for all bodies is same and is equal to the emissive power of a perfectly black body.

Ans: e

63. All radiations in a black body are

- (a) reflected
- (b) refracted
- (c) transmitted
- (d) absorbed
- (e) partly reflected and partly absorbed.

Ans: d

64. According to Kirchoff's law, the ratio of emissive power to absorptivity for all bodies is equal to the emissive power of a

- (a) grey body
- (b) brilliant white polished body
- (c) red hot body
- (d) black body
- (e) none of the above.

Ans: d

65. The concept of overall coefficient of heat transfer is used in case of heat transfer by

- (a) conduction
- (b) convection
- (c) radiation
- (d) conduction and convection
- (e) convection and radiation.

Ans: d

66. The unit of overall coefficient of heat transfer is

- (a) kcal/m²
- (b) kcal/hr °C
- (c) kcal/m² hr °C
- (4) kcal/m hr °C
- (e) kcal/m³ hr °C.

Ans: c

68. Joule sec is the unit of

- (a) universal gas constant
- (b) kinematic viscosity
- (c) thermal conductivity
- (d) Planck's constant
- (e) none of the above.

Ans: d

69. The value of Prandtl number for air is about

- (a) 0.1
- (b) 0.3
- (c) 0.7
- (d) 1.7
- (e) 10.5.

Ans: c

70. The value of the wavelength for maximum emissive power is given by —

- (a) Wien's law
- (b) Planck's law
- (c) Stefan's law
- (d) Fourier's law
- (e) Kirchhoff's law.

Ans: a

72. Log mean temperature difference in case of counter flow compared to parallel flow will be

- (a) same
- (b) more
- (c) less
- (d) depends on other factors
- (e) none of the above.

Ans: b

73. The energy distribution of an ideal reflector at higher temperatures is largely in the range of

- (a) shorter wavelength
- (b) longer wavelength
- (c) remains same at all wavelengths
- (d) wavelength has nothing to do with it
- (e) none of the above.

Ans: a

74. Total emissivity of polished silver compared to black body is

- (a) same
- (b) higher
- (c) more or less same
- (d) very much lower
- (e) very much higher.

Ans: d

75. According to Stefan-Boltzmann law, ideal radiators emit radiant energy at a rate proportional to

- (a) absolute temperature
- (b) square of temperature
- (c) fourth power of absolute temperature
- (d) fourth power of temperature
- (e) cube of absolute temperature.

Ans: c

76. Which of the following property of air does not increase with rise in temperature

- (a) thermal conductivity
- (b) thermal diffusivity
- (c) density
- (d) dynamic viscosity
- (e) kinematic viscosity.

Ans: c

77. The unit of Stefan Boltzmann constant is

- (a) watt/cm² °K
- (b) watt/cm⁴ °K
- (c) watt²/cm °K⁴
- (d) watt/cm² °K⁴
- (e) watt/cm² °K².

Ans: d

78. In free con-vection heat transfer, Nusselt number is function of

- (a) Grashoff no. and Reynold no.
- (b) Grashoff no. and Prandtl no.
- (c) Prandtl no. and Reynold no.
- (d) Grashoff no., Prandtl no. and Reynold no.
- (e) none of the above.

Ans: b

79. Stefan Boltzmann law is applicable for heat transfer by

- (a) conduction
- (b) convection
- (c) radiation
- (d) conduction and radiation combined
- (e) convection and radiation combined.

Ans: c

80. The thermal diffusivities for gases are generally

- (a) more than those for liquids
- (b) less than those for liquids
- (c) more than those for solids
- (d) dependent on the viscosity
- (e) same as for the liquids.

Ans: a

81. The thermal diffusivities for solids are generally

- (a) less than those for gases
- (b) less than those for liquids
- (c) more than those for liquids and gases
- (d) more or less same as for liquids and gases
- (e) zero.

Ans: c

83. Thermal diffusivity of a substance is

- (a) directly proportional to thermal conductivity
- (b) inversely proportional to density of substance
- (c) inversely proportional to specific heat
- (d) all of the above
- (e) none of the above.

Ans: d

85. The ratio of the emissive power and absorptive power of all bodies is the same and is equal to the emissive power of a perfectly black body. This statement is known as

- (a) Kirchhoff's law
- (b) Stefan's law
- (c) Wien's law
- (d) Planck's law
- (e) Black body law.

Ans: a

86. According to Stefan's law, the total radiation from a black body per second per unit area is proportional to

- (a) absolute temperature
- (b) T^2
- (c) T^5
- (d) t
- (e) $1/T$.

Ans: d

87. According to Wien's law, the wavelength corresponding to maximum energy is proportion to

- (a) absolute temperature (T)
- (b) $1/T$
- (c) f
- (d) t
- (e) $1/\lambda$.

Ans: a

88. Depending on the radiating properties, a body will be white when

- (a) $p = 0, x = 0$ and $a = 1$
- (b) $p = 1, T = 0$ and $a = 0$
- (c) $p = 0, x = 1$ and $a = 0$
- (d) $x = 0, a + p = 1$
- (e) $a = 0, x + p = 1$.

where $a =$ absorptivity, $p =$ reflectivity, $x =$ transmissivity

Ans: b

89. Depending on the radiating properties, a body will be black when

- (a) $p = 0, x = 0$ and $a = 1$
- (b) $p = 1, T = 0$ and $a = 0$
- (c) $p = 0, x = 1$ and $a = 0$
- (d) $x = 0, a + p = 0$
- (e) $a = 0, x + p = 1$.

where $a =$ absorptivity, $p =$ reflectivity, $X =$ transmissivity.

Ans: a

90. Depending on the radiating properties, a body will be opaque when

- (a) $p = 0, x = 0$ and $a = 1$
- (b) $p = 1, x = 0$ and $a = 0$
- (c) $p = 0, x = 1$ and $a = 0$
- (d) $x = 0, a + p = 1$
- (e) $a = 0, x + p = 1$.

where $a =$ absorptivity, $p =$ reflectivity, $X =$ transmissivity.

Ans: d

91. The total emissivity power is defined as the total amount of radiation emitted by a black body per unit

- (a) temperature
- (b) thickness
- (c) area
- (d) time
- (e) area and time.

Ans: d

92. The ratio of the energy absorbed by the body to total energy falling on it is called

- (a) absorptive power
- (b) emissive power
- (c) absorptivity
- (d) emissivity
- (e) none of the above.

Ans: a

93. 40% of incident radiant energy on the surface of a thermally transparent body is reflected back. If the transmissivity of the body be 0.15, then the emissivity of surface is

- (a) 0.45
- (b) 0.55
- (c) 0.40
- (d) 0.75

(e) 0.60.

Ans: a

94. The amount of radiation mainly depends on

- (a) nature of body
- (b) temperature of body
- (c) type of surface of body
- (d) all of the above
- (e) none of the above.

Ans: d

95. The emissive power of a body depends upon its

- (a) temperature
- (b) wave length
- (c) physical nature
- (d) all of the above
- (e) none of the above.

Ans: d

96. Two plates spaced 150 mm apart are maintained at 1000°C and 70°C. The heat transfer will take place mainly by

- (a) convection
- (b) free convection
- (c) forced convection
- (d) radiation
- (e) radiation and convection.

Ans: d

97. Absorptivity of a body will be equal to its emissivity

- (a) at all temperatures
- (b) at one particular temperature
- (c) when system is under thermal equilibrium
- (d) at critical temperature
- (e) for a polished body.

Ans: c

98. In regenerator type heat exchanger, heat transfer takes place by

- (a) direct mixing of hot and cold fluids
- (b) a complete separation between hot and cold fluids
- (c) flow of hot and cold fluids alternately over a surface
- (d) generation of heat again and again
- (e) indirect transfer.

Ans: c

99. A perfect black body is one which

- (a) is black in colour
- (b) reflects all heat
- (c) transmits all heat radiations
- (d) absorbs heat radiations of all wave lengths falling on it
- (e) fully opaque.

Ans: d

100. Planck's law holds good for

- (a) black bodies
- (b) polished bodies
- (c) all coloured bodies
- (d) all of the above
- (e) none of the above.

Ans: a

101. If the temperature of a solid surface changes from 27°C to 627°C , then its emissive power changes in the ratio of

- (a) 3
- (b) 6
- (c) 9
- (d) 27
- (e) 81.

Ans: e

102. Depending on the radiating properties, body will be transparent when

- (a) $p = 0, x = 0$ and $a = 1$
- (b) $p=1, x = 0, \text{ and } a = 0$
- (c) $p = 0, T= 1, \text{ and } a = 0$
- (d) $X = 0, a + p = 1$
- (e) $a = 0, x + p = 1.$

Ans: c

103. A grey body is one whose absorptivity

- (a) varies with temperature
- (b) varies with the wave length of incident ray
- (c) varies with both
- (d) does not vary with temperature and wave length of the incident ray
- (e) there is no such criterion.

Ans: d